

# Child SCOAT6

Gavin A Davis <sup>1,2</sup>, Jon S Patricios <sup>3</sup>, Laura K Purcell <sup>4</sup>, Vicki Anderson <sup>5,6</sup>, Gerry Gioia <sup>7</sup>, Christopher C Giza <sup>8,9</sup>, Keith Owen Yeates <sup>10</sup>, Osman Hassan Ahmed <sup>11,12</sup>, Cheri Blauwet <sup>13</sup>, Daniel Corwin <sup>14</sup>, Christina L Master <sup>15</sup>, Geoff Schneider <sup>16</sup>, Jacqueline van Ierssel <sup>17</sup>, Ruben J Echemendia <sup>18,19</sup>, Pierre Fremont <sup>20</sup>, Gordon Ward Fuller <sup>21</sup>, Kimberly G Harmon <sup>22</sup>, Stanley A Herring <sup>23</sup>, Kirsten Holte <sup>24</sup>, Mike Loosemore <sup>25</sup>, Michael Makdissi <sup>26,27</sup>, Michael McCrea <sup>28</sup>, William P Meehan, III <sup>29,30</sup>, Patrick O'Halloran <sup>31</sup>, Zahra Premji <sup>32</sup>, Margot Putukian <sup>33</sup>, Isla Jordan Shill <sup>34</sup>, Michael Turner <sup>35</sup>, Kenzie Vaandering <sup>36</sup>, Nick Webborn <sup>37</sup>, Kathryn J Schneider <sup>38</sup>

## child SCOAT6<sup>TM</sup>

### Sport Concussion Office Assessment Tool

#### For Children Ages 8 to 12 Years

#### What is the Child SCOAT6?\*

The Child SCOAT6 is a tool for evaluating concussions in a controlled office environment by Health Care Professionals (HCP) typically from 72 hours (3 days) following a sport-related concussion.

The diagnosis of concussion is a clinical determination made by an HCP. The various components of the Child SCOAT6 may assist with the clinical assessment and help guide individualised management.

The Child SCOAT6 is used for evaluating athletes aged 8 - 12 years. For athletes aged 13 years and older, please use the SCOAT6.

Brief verbal instructions for some components of the Child SCOAT6 are included. Detailed instructions for use of the Child SCOAT6 are provided in an accompanying document. Please read through these instructions carefully before using the Child SCOAT6.

This tool may be freely copied in its current form for distribution to individuals, teams, groups, and organisations.

Any alteration (including translations and digital re-formatting), re-branding, or sale for commercial gain is not permissible without the expressed written consent of BMJ and the Concussion in Sport Group (CISG).

#### Completion Guide

Blue: Complete only at first assessment

Green: Recommended part of assessment

Orange: Optional part of assessment

Athlete's Name: <input type="text"/>		
Date of Birth: <input type="text"/>	Sex: Male <input type="checkbox"/> Female <input type="checkbox"/> Prefer Not To Say <input type="checkbox"/>	
Sport: <input type="text"/>		
Age First Played Contact Sport: <input type="text"/>	School Class/Grade/Level: <input type="text"/>	
Handedness (Writing): L <input type="checkbox"/> R <input type="checkbox"/> Ambidextrous <input type="checkbox"/>	Handedness (Sport): L <input type="checkbox"/> R <input type="checkbox"/> Ambidextrous <input type="checkbox"/>	
Dominant Leg (Sport): L <input type="checkbox"/> R <input type="checkbox"/> Ambidextrous <input type="checkbox"/>		
Name of Accompanying Parent/Carer: <input type="text"/>		
Examiner: <input type="text"/>	Date of Examination: <input type="text"/>	
Referring Physician's Name: <input type="text"/>		
Referring Physician's Contact Details: <input type="text"/>		

\* In reviewing studies informing the SCOAT6 and Child SCOAT6, the period defined for the included papers was 3–30 days. HCPs may choose to use the Child SCOAT6 beyond this timeframe but should be aware of the parameters of the review.

For use by Health Care Professionals Only

Child SCOAT6<sup>TM</sup>

Developed by: The Concussion in Sport Group (CISG)

Supported by:



International  
Olympic  
Committee



**Correspondence to** Professor Gavin A Davis,  
Murdoch Children's Research Institute, Parkville VIC  
3052, Victoria, Australia; gavin.davis@me.com


**child SCOAT6™**
**Sport Concussion Office Assessment Tool**  
 For Children Ages 8 to 12 Years


### Current Injury

 Removal From Play: Immediate ☐ Continued to play for \_\_\_\_\_ mins ☐  
 Walked off ☐ Assisted off ☐ Stretchered off ☐

 Date of Injury: 
**Description** - include mechanism of injury, presentation, management since the time of injury and trajectory of care since injury:

 Date Symptoms First Appeared: 

 Date Symptoms First Reported: 

### History of Head Injuries

Date/Year	Description - include mechanism of injury, main symptoms, recovery time	Management - including time off school or sport
<input type="text"/>	<input type="text"/>	<input type="text"/>

### History of Any Neurological, Psychological, Psychiatric or Learning Disorders

Diagnosis	Year Diagnosed	Management Including Medication
<input type="checkbox"/> Migraine	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Chronic headache	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Depression	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Anxiety	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Syncope	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Epilepsy/seizures	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Attention deficit hyper-activity disorder (ADHD)	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Learning disorder/ dyslexia	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Developmental Co-ordination Disorder	<input type="text"/>	<input type="text"/>
<input type="checkbox"/> Other _____	<input type="text"/>	<input type="text"/>



## Item

## Dose

Frequency

Reason Taken

## Family Member

## Diagnosis

## Management Including Medication

☐ Depression

#### **Anxiety**

☐ Attention deficit hyperactivity disorder (ADHD)

☐ Learning disorder/  
dyslexia

#### **Migraine**

☐ Other \_\_\_\_\_

**Additional Notes:**



## Child Report

Child to complete all 3 symptom boxes

### Box 1

Symptom	Not at all/never	A little/rarely	Somewhat/sometimes	A lot/often
I have headaches	0	1	2	3
I feel dizzy	0	1	2	3
I feel like the room is spinning	0	1	2	3
I feel like I'm going to faint	0	1	2	3
Things are blurry when I look at them	0	1	2	3
I see double	0	1	2	3
I feel sick to my stomach	0	1	2	3
I get tired a lot	0	1	2	3
I get tired easily	0	1	2	3
I have trouble paying attention	0	1	2	3
I get distracted easily	0	1	2	3
I have a hard time concentrating	0	1	2	3
I have problems remembering what people tell me	0	1	2	3
I have problems following directions	0	1	2	3
I daydream too much	0	1	2	3
I get confused	0	1	2	3
I forget things	0	1	2	3
I have problems finishing things	0	1	2	3
I have trouble figuring things out	0	1	2	3
It's hard for me to learn new things	0	1	2	3

Box 1: Total Number of Symptoms:  of 20      Symptom Severity Score:  of 60

### Box 2

Symptom	Not at all/never	A little/rarely	Somewhat/sometimes	A lot/often
My neck hurts	0	1	2	3
I have problems with bright lights	0	1	2	3
I have problems with loud noise	0	1	2	3
I feel sleepy or drowsy	0	1	2	3
I am sleeping more than usual	0	1	2	3
I have difficulty falling asleep or staying asleep at night	0	1	2	3
I have problems with balance	0	1	2	3
I am thinking more slowly	0	1	2	3
I am more emotional	0	1	2	3
Things annoy me easily	0	1	2	3
I am sad	0	1	2	3
I have problems looking up at the board after looking at work on my desk	0	1	2	3

Box 2: Total Number of Symptoms:  of 12      Symptom Severity Score:  of 36





## Child Report (Continued)

### Box 3

Do the symptoms get worse with physical activity?	Y	N
Do the symptoms get worse with trying to think?	Y	N

Overall rating for child to answer:

On a scale of 0 to 10 (where 10 is normal), how do you feel now? Very Bad 0 1 2 3 4 5 6 7 8 9 10 Very Good

If not 10, in what way do you feel different?

### Child Report (Box 1 + Box 2)

Total Number of Symptoms:  of 32

Symptom Severity Score:  of 96

## Parent Report

Parent to complete all 3 symptom boxes

### Box 1

The Child...

Symptom	Not at all/never	A little/rarely	Somewhat/sometimes	A lot/often
has headaches	0	1	2	3
feels dizzy	0	1	2	3
has a feeling that the room is spinning	0	1	2	3
feels faint	0	1	2	3
has blurred vision	0	1	2	3
has double vision	0	1	2	3
experiences nausea	0	1	2	3
gets tired a lot	0	1	2	3
gets tired easily	0	1	2	3
has trouble sustaining attention	0	1	2	3
is distracted easily	0	1	2	3
has difficulty concentrating	0	1	2	3
has problems remembering what he/she is told	0	1	2	3
has difficulty following directions	0	1	2	3
tends to daydream	0	1	2	3
gets confused	0	1	2	3
is forgetful	0	1	2	3
has difficulty completing tasks	0	1	2	3
has poor problem-solving skills	0	1	2	3
has problems learning	0	1	2	3

Box 1: Total Number of Symptoms:  of 20

Symptom Severity Score:  of 60



## Parent Report (Continued)

### Box 2

The Child...

Symptom	Not at all/never	A little/rarely	Somewhat/sometimes	A lot/often
has a sore neck	0	1	2	3
is sensitive to light	0	1	2	3
is sensitive to noise	0	1	2	3
appears drowsy	0	1	2	3
is sleeping more than usual	0	1	2	3
has difficulty falling asleep or staying asleep at night	0	1	2	3
has balance problems	0	1	2	3
is thinking more slowly	0	1	2	3
acts more emotional	0	1	2	3
acts irritable	0	1	2	3
appears sad	0	1	2	3
has difficulty shifting vision in the classroom (i.e. looking from work on a desk to board)	0	1	2	3

Box 2: Total Number of Symptoms:  of 12      Symptom Severity Score:  of 36

### Box 3

Do the symptoms get worse with physical activity?	Y	N
Do the symptoms get worse with trying to think?	Y	N

Overall rating for parent/teacher/coach/carer to answer:

On a scale of 0 to 100% (where 100% is normal), how would you rate the child now?

If not 100%, in what way does the child seem different?

### Parent Report (Box 1 + Box 2)

Total Number of Symptoms:  of 32      Symptom Severity Score:  of 96

## PACE Self-Efficacy Questionnaire - Self Report

A measure that indicates the degree of the child's confidence in their actions affecting recovery.

Questionnaire contained in Child SCOAT6 Supplementary Material



## Verbal Cognitive Tests

### Immediate Memory

All 3 trials must be administered irrespective of the number correct on Trial 1. Administer at the rate of one word per second in a monotone voice.

**Trial 1:** Say *"I am going to test your memory. I will read you a list of words and when I am done, repeat back as many words as you can remember, in any order."*

**Trials 2 and 3:** Say *"I am going to repeat the same list. Repeat back as many words as you can remember in any order, even if you said the word before in a previous trial."*

Word list used: A ☐ B ☐ C ☐

				Alternate Lists	
List A	Trial 1	Trial 2	Trial 3	List B	List C
Jacket	0 1	0 1	0 1	Finger	Baby
Arrow	0 1	0 1	0 1	Penny	Monkey
Pepper	0 1	0 1	0 1	Blanket	Perfume
Cotton	0 1	0 1	0 1	Lemon	Sunset
Movie	0 1	0 1	0 1	Insect	Iron
Dollar	0 1	0 1	0 1	Candle	Elbow
Honey	0 1	0 1	0 1	Paper	Apple
Mirror	0 1	0 1	0 1	Sugar	Carpet
Saddle	0 1	0 1	0 1	Sandwich	Saddle
Anchor	0 1	0 1	0 1	Wagon	Bubble
<b>Trial Total</b>					

Immediate Memory Total \_\_\_\_\_ of 30

Time last trial completed: \_\_\_\_\_

## Digits Backwards

Administer at the rate of one word per second in a monotone voice.

**Say** *"I am going to read a string of numbers and when I am done, you repeat them back to me in reverse order of how I read them to you. For example, if I say 7-1, you would say 1-7. So, if I said 6-8 you would say? (8-6)"*

Digit list used: A ☐ B ☐ C ☐

List A	List B	List C	Y	N	0	1
2-7	9-2	7-8	Y	N	0	1
5-9	6-1	5-1	Y	N	0	1
7-8-2	3-8-2	2-7-1	Y	N	0	1
9-2-6	5-1-8	4-7-9	Y	N	0	1
4-1-8-3	2-7-9-3	1-6-8-3	Y	N	0	1
9-7-2-3	2-1-6-9	3-9-2-4	Y	N	0	1
1-7-9-2-6	4-1-8-6-9	2-4-7-5-8	Y	N	0	1
4-1-7-5-2	9-4-1-7-5	8-3-9-6-4	Y	N	0	1
6-0-1-3-5-7	2-5-1-3-9-8	0-7-5-8-1-6	Y	N	0	1
6-1-2-8-0-7	0-8-5-1-9-4	0-2-8-4-7-1	Y	N	0	1
				<b>Digits score</b>		
				<b>of 4</b>		

## Days in Reverse Order

**Say** *"Now tell me the days of the week in reverse order. Start with the last day and go backward. So you'll say Sunday, Saturday, and so on... Go ahead."* Start stopwatch and CIRCLE each correct response:

Sunday Saturday Friday Thursday Wednesday Tuesday Monday

Time Taken to Complete (secs): \_\_\_\_\_ (N < 30 sec) Number of Errors: \_\_\_\_\_



## Symbol Digit Modalities Test

A measure of psychomotor processing speed.

If clinically indicated based on symptoms and clinical findings

**SDMT contained in Child SCOAT6 Supplementary Material**

## Examination

### Orthostatic Vital Signs

Take the child's blood pressure and pulse via digital sphygmomanometer after lying supine for 2 minutes; and then again after standing unsupported for 2 minutes. An option is to perform an additional assessment between lying and standing: after sitting upright for 2 minutes. The child is asked if they experience any symptoms such as: dizziness or light-headedness, fainting, blurred or fading vision, nausea, fatigue, or lack of concentration.

Orthostatic Vital Signs	Supine (after 2 minutes)	Standing (after 2 minutes)
Blood Pressure (mmHg)		
Heart Rate (bpm)		
Symptoms <sup>1</sup> • Dizziness or light-headedness • Fainting • Blurred or fading vision • Nausea • Fatigue • Lack of concentration	No <input type="checkbox"/> Yes <input type="checkbox"/> If yes: Description	No <input type="checkbox"/> Yes <input type="checkbox"/> If yes: Description
Results	<input type="checkbox"/> Normal <input type="checkbox"/> Abnormal	

Orthostatic hypotension: a drop in systolic BP  $\geq 20$  mmHg between supine and standing positions. Orthostatic tachycardia: an elevation in HR of  $\geq 30$  bpm when transitioning between the supine and standing positions, in the absence of orthostatic hypotension.

### Cervical Spine Assessment

Cervical Spine Palpation	Signs and Symptoms		Location
Muscle Spasm	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Midline Tenderness	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Paravertebral Tenderness	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Cervical Active Range of Motion	Result		
Flexion (50-80°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Extension (45-95°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Right Lateral Flexion (30-55°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Left Lateral Flexion (30-55°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Right Rotation (50-90°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	
Left Rotation (50-90°)	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal	

Notes:



## Neurological Examination

### Cranial Nerves

Normal ☐ Abnormal ☐ Not tested ☐

Notes:

### Finger to Nose

Eyes Open:

Left Hand: Normal ☐ Abnormal ☐ Not tested ☐

Right Hand: Normal ☐ Abnormal ☐ Not tested ☐

Eyes Closed:

Left Hand: Normal ☐ Abnormal ☐ Not tested ☐

Right Hand: Normal ☐ Abnormal ☐ Not tested ☐

### Other Neurological Findings

Limb Tone: Normal ☐ Abnormal ☐ Not tested ☐

Strength: Normal ☐ Abnormal ☐ Not tested ☐

Deep Tendon Reflexes: Normal ☐ Abnormal ☐ Not tested ☐

Sensation: Normal ☐ Abnormal ☐ Not tested ☐

Cerebellar Function: Normal ☐ Abnormal ☐ Not tested ☐

Comments:

## Balance

Barefoot on a firm surface with or without foam mat

Foot Tested: Left ☐ Right ☐ (i.e. test the **non-dominant** foot)

### Modified BESS

Double Leg Stance:  of 10

Tandem Stance:  of 10

Single Leg Stance:  of 10

Total Errors:  of 30

### On Foam

Double Leg Stance:  of 10

Tandem Stance:  of 10

Single Leg Stance:  of 10

Total Errors:  of 30



### Timed Tandem Gait

Place a 3-metre-long line on the floor/firm surface with athletic tape. The task should be timed.

Say *"Please walk heel-to-toe quickly to the end of the tape, turn around and come back as fast as you can without separating your feet or stepping off the line."*

Time to Complete Tandem Gait Walking (seconds)				
Trial 1	Trial 2	Trial 3	Average 3 Trials	Fastest Trial

Abnormal/failed to complete ☐ Unstable/sway ☐ Fall/over-step ☐ Dizzy/nauseated ☐

### Complex Tandem Gait

#### Forward

Say *"Please walk heel-to-toe quickly five steps forward, then continue forward with eyes closed five steps"*

1 point for each step off the line, 1 point for truncal sway.

Forward Eyes Open Points:

Forward Eyes Closed Points:

Forward Total Points:

#### Backward

Say *"Please walk heel-to-toe again, backwards five steps eyes open, then continue backwards five steps with eyes closed."* 1 point for each step off the line, 1 point for truncal sway.

Backward Eyes Open Points:

Backward Eyes Closed Points:

Backward Total Points:

Total Points (Forward + Backward):

### Dual Task Gait

Only perform if child successfully completes Complex Tandem Gait

Say *"Now, while you are walking heel-to-toe, I will ask you to count backwards out loud by 7s (or 3s) / recite the months of the year (or days of the week) in reverse order"* (select one cognitive task). Allow for a verbal practice attempt of the cognitive task selected.

Cognitive Tasks												
Trial 1	95	88	81	74	67	60	53	46				
(Subtract serial 7s)												
OR												
(Subtract serial 3s)	97	94	91	88	85	82	79	76				
OR Trial 2	December	November	October	September	August	July	June	May	April	March	February	January
(Months backward)												
OR												
(Days backward)	Thursday	Wednesday	Tuesday	Monday	Sunday	Saturday	Friday					

Before attempting the dual task: *"Good. Now I will ask you to walk heel-to-toe calling the answers out loud at the same time. Are you ready?"*

Cognitive Accuracy: Number Correct:  Number Incorrect:  Average Time (s):

Comments:





## Visio-Vestibular Examination

### Smooth Pursuits

Patient-reported Symptom Provocation:

Worsening Headache: Yes ☐ No ☐ Dizziness: Yes ☐ No ☐  
 Eye Fatigue: Yes ☐ No ☐ Eye Pain: Yes ☐ No ☐ Nausea: Yes ☐ No ☐

Or Physical Signs:

Jerky or Jumpy Eye Movements: Yes ☐ No ☐ >3 Beats of Nystagmus: Yes ☐ No ☐

### Fast Saccades

Horizontal Saccades:

Worsening Headache: Yes ☐ No ☐ Dizziness: Yes ☐ No ☐  
 Eye Fatigue: Yes ☐ No ☐ Eye Pain: Yes ☐ No ☐ Nausea: Yes ☐ No ☐

Vertical Saccades:

Worsening Headache: Yes ☐ No ☐ Dizziness: Yes ☐ No ☐  
 Eye Fatigue: Yes ☐ No ☐ Eye Pain: Yes ☐ No ☐ Nausea: Yes ☐ No ☐

### Gaze Stability Testing (The Angular Vestibular-Ocular Reflex)

Vertical Gaze Stability:

Worsening Headache: Yes ☐ No ☐ Dizziness: Yes ☐ No ☐  
 Eye Fatigue: Yes ☐ No ☐ Eye Pain: Yes ☐ No ☐ Nausea: Yes ☐ No ☐

Horizontal Gaze Stability:

Worsening Headache: Yes ☐ No ☐ Dizziness: Yes ☐ No ☐  
 Eye Fatigue: Yes ☐ No ☐ Eye Pain: Yes ☐ No ☐ Nausea: Yes ☐ No ☐

### Near Point of Convergence Testing

Distance:  cm

### Left and Right Monocular Accommodation

Left Eye Distance:  cm Right Eye Distance:  cm

### Complex Tandem Gait (if not tested in Balance)

Complex Tandem Gait Score:

## Pediatric Athlete Mental Health

### Pediatric Anxiety – Short Form 8a

If clinically indicated based on symptoms and clinical findings

Pediatric Anxiety Questionnaire contained in Child SCOAT6 Supplementary Material

### Pediatric Depressive Symptoms – Short Form 8a

If clinically indicated based on symptoms and clinical findings

Pediatric Depressive Questionnaire contained in Child SCOAT6 Supplementary Material





## Pediatric Athlete Mental Health (Continued)

### Pediatric Sleep Disturbance – Short Form 4a

If clinically indicated based on symptoms and clinical findings

**Pediatric Sleep Disturbance Questionnaire contained in Child SCOAT6 Supplementary Material**

### Pediatric Sleep-Related Impairment – Short Form 4a

If clinically indicated based on symptoms and clinical findings

**Pediatric Sleep-Related Impairment Questionnaire contained in Child SCOAT6 Supplementary Material**

### The Pediatric Fear Avoidance Behavior after Traumatic Brain Injury Questionnaire (PFAB-TBI)

A measure to identify fear avoidance behaviour, which may contribute to poorer outcomes/persisting symptoms post concussion, which may benefit from psychological intervention.

**PFAB-TBI Questionnaire contained in Child SCOAT6 Supplementary Material**

## Delayed Word Recall

Minimum of 5 minutes after immediate recall

Say *"Do you remember that list of words I read a few times earlier? Tell me as many words from the list as you can remember in any order."*

Word list used:	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	Alternate Lists	
List A	Score			List B	List C
Jacket	0 1			Finger	Baby
Arrow	0 1			Penny	Monkey
Pepper	0 1			Blanket	Perfume
Cotton	0 1			Lemon	Sunset
Movie	0 1			Insect	Iron
Dollar	0 1			Candle	Elbow
Honey	0 1			Paper	Apple
Mirror	0 1			Sugar	Carpet
Saddle	0 1			Sandwich	Saddle
Anchor	0 1			Wagon	Bubble

Score:  of 10

Record Actual Time (mins) Since Completing Immediate Recall:

## Computerised Cognitive Test Results (if used)

Not Done ☐

Test Battery Used:

Recent Baseline - if performed (Date):

Post-Injury Result (Rest):

Post-Injury Result (Post-Exercise Stress):

## Graded Aerobic Exercise Test

Not Done ☐

Exclude contra-indications: cardiac condition, respiratory disease, significant vestibular symptoms, motor dysfunction, lower limb injuries, cervical spine injury.

Protocol Used:



## Overall Assessment

## Summary:

## Management and Follow-up Plan

## Recommendations regarding return to:

School/Class:

Sport:

Assessment by:

Name:

☐ Athletic Trainer/Therapist

☐ Exercise Physiologist

☐ Neurologist

☐ Neuropsychologist

☐ Neurosurgeon

☐ Ophthalmologist

☐ Optometrist

☐ Paediatrician

☐ Physiatrist/Rehab Phys

☐ Physiotherapist

☐ Psychologist

☐ Psychiatrist

☐ Sport and Exercise Medicine Phys

☐ Other

 Neuroimaging: Not Required ☐ Required and Requested ☐ Already Performed and Images Reviewed ☐

Details:

Brain: CT ☐ MRI ☐Cervical Spine: XR ☐ CT ☐ MRI ☐ Other ☐


Details:

## Pharmacotherapy Prescribed:

Date of Review:

Date of Follow-up:



### Additional Clinical Notes

### Return-to-Learn (RTL) Strategy

Facilitating RTL is a vital part of the recovery process for student-athletes. HCPs should work with stakeholders on education and school policies to facilitate academic support, including accommodations/learning adjustments for students with SRC when needed. Academic support should address risk factors for greater RTL duration (e.g., social determinants of health, higher symptom burden) by adjusting environmental, physical, curricular, and testing factors as needed. **Not all athletes will need a RTL strategy or academic support.** If symptom exacerbation occurs during cognitive activity or screen time, or difficulties with reading, concentration, or memory or other aspects of learning are reported, clinicians should consider implementation of a RTL strategy at the time of diagnosis and during the recovery process. When the RTL strategy is implemented, it can begin following an initial period of relative rest (Stage 1: 24-48 hrs), with an incremental increase in cognitive load (Stages 2 to 4). Progression through the strategy is symptom limited (i.e., no more than a mild exacerbation of current symptoms related to the current concussion) and its course may vary across individuals based on tolerance and symptom resolution. Further, while the RTL and RTS strategies can occur in parallel, student-athletes should complete full RTL before unrestricted RTS.

Step	Mental Activity	Activity at Each Step	Goal
1	Daily activities that do not result in more than a mild exacerbation* of symptoms related to the current concussion.	Typical activities during the day (e.g., reading) while minimizing screen time. Start with 5–15 min at a time and increase gradually.	Gradual return to typical activities.
2	School activities.	Homework, reading, or other cognitive activities outside of the classroom.	Increase tolerance to cognitive work.
3	Return to school part time.	Gradual introduction of schoolwork. May need to start with a partial school day or with greater access to rest breaks during the day.	Increase academic activities.
4	Return to school full time.	Gradually progress school activities until a full day can be tolerated without more than mild* symptom exacerbation.	Return to full academic activities and catch up on missed work.

**NOTE:** Following an initial period of relative rest (24-48 hours following injury at Step 1), athletes can begin a gradual and incremental increase in their cognitive load. Progression through the strategy for students should be slowed when there is more than a mild and brief symptom exacerbation.

\*Mild and brief exacerbation of symptoms is defined as an increase of no more than 2 points on a 0-10 point scale (with 0 representing no symptoms and 10 the worst symptoms imaginable) for less than an hour when compared with the baseline value reported prior to cognitive activity.



## Return-to-Sport (RTS) Strategy

Return to sport participation after an SRC follows a graduated stepwise strategy, an example of which is outlined in Table 2. RTS occurs in conjunction with return to learn (see RTL strategy) and under the supervision of a qualified HCP. Following an initial period of relative rest (step 1: approximately 24-48 hours), clinicians can implement step 2 [i.e., light (step 2A) and then moderate (step 2B) aerobic activity] of the RTS strategy as a treatment of acute concussion. The athlete may then advance to steps 3-6 on a time course dictated by symptoms, cognitive function, clinical findings, and clinical judgement. Differentiating early activity (step 1), aerobic exercise (step 2), and individual sport-specific exercise (step 3) as part of the treatment of SRC from the remainder of the RTS progression (steps 4-6) can be useful for the athlete and their support network (e.g., parents, coaches, administrators, agents). Athletes may be moved into the later stages that involve risk of head impact (steps 4-6 and step 3 if there is any risk of head impact with sport-specific activity) of the RTS strategy following authorization by the HCP and after resolution of any new symptoms, abnormalities in cognitive function, and clinical findings related to the current concussion. Each step typically takes at least 24 hours. Clinicians and athletes can expect a minimum of 1 week to complete the full rehabilitation strategy, but typical unrestricted RTS can take up to one month post-SRC. The time frame for RTS may vary based on individual characteristics, necessitating an individualized approach to clinical management. Athletes having difficulty progressing through the RTS strategy or with symptoms and signs that are not progressively recovering beyond the first 2-4 weeks may benefit from rehabilitation and/or involvement of a multidisciplinary team of HCP experienced in managing SRC. Medical determination of readiness to return to at-risk activities should occur prior to returning to any activities at risk of contact, collision or fall (e.g. multiplayer training drills), which may be required prior to any of steps 3-6, depending on the nature of the sport or activity that the athlete is returning to and in keeping with local laws/requirements.

Step	Exercise Strategy	Activity at Each Step	Goal
1	Symptom-limited activity.	Daily activities that do not exacerbate symptoms (e.g., walking).	Gradual reintroduction of work/school.
2	Aerobic exercise <b>2A – Light</b> (up to approx. 55% max HR) <b>then</b> <b>2B – Moderate</b> (up to approximately 70% max HR)	Stationary cycling or walking at slow to medium pace. May start light resistance training that does not result in more than mild and brief exacerbation* of concussion symptoms.	Increase heart rate.
3	Individual sport-specific exercise <b>NOTE:</b> if sport-specific exercise involves any risk of head impact, medical determination of readiness should occur prior to step 3.	Sport-specific training away from the team environment (e.g., running, change of direction and/or individual training drills away from the team environment). No activities at risk of head impact.	Add movement, change of direction.
Steps 4-6 should begin after resolution of any symptoms, abnormalities in cognitive function, and any other clinical findings related to the current concussion, including with and after physical exertion.			
4	Non-contact training drills.	Exercise to high intensity including more challenging training drills (e.g., passing drills, multiplayer training). Can integrate into team environment.	Resume usual intensity of exercise, coordination, and increased thinking.
5	Full contact practice.	Participate in normal training activities.	Restore confidence and assess functional skills by coaching staff.
6	Return to sport.	Normal game play.	

maxHR = predicted maximal Heart Rate according to age (i.e., 220-age)

Age Predicted Maximal HR= 220-age	Mild Aerobic Exercise	Moderate Aerobic Exercise
55%	220-age x 0.55 = training target HR	
70%		220-age x 0.70 = training target HR

**NOTE:** \*Mild and brief exacerbation of symptoms (i.e., an increase of no more than 2 points on a 0-10 point scale for less than an hour when compared with the baseline value reported prior to physical activity). Athletes may begin Step 1 (i.e., symptom-limited activity) within 24 hours of injury, with progression through each subsequent step typically taking a minimum of 24 hours. If more than mild exacerbation of symptoms (i.e., more than 2 points on a 0-10 scale) occurs during Steps 1 -3, the athlete should stop and attempt to exercise the next day. If an athlete experiences concussion-related symptoms during Steps 4-6, they should return to Step 3 to establish full resolution of symptoms with exertion before engaging in at-risk activities. Written determination of readiness to RTS should be provided by an HCP before unrestricted RTS as directed by local laws and/or sporting regulations.



**Author affiliations**

- <sup>1</sup>Murdoch Children's Research Institute, Parkville, Victoria, Australia
- <sup>2</sup>Cabrini Health, Malvern, Victoria, Australia
- <sup>3</sup>Wits Sport and Health (WiSH), School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg-Braamfontein, South Africa
- <sup>4</sup>Department of Pediatrics, McMaster University, Hamilton, Ontario, Canada
- <sup>5</sup>Child Neuropsychology, Murdoch Children's Research Institute, Parkville, Victoria, Australia
- <sup>6</sup>Psychology, University of Melbourne, Parkville, Victoria, Australia
- <sup>7</sup>Children's National Health System, Washington, District of Columbia, USA
- <sup>8</sup>Neurosurgery, UCLA Steve Tisch BrainSPORT Program, Los Angeles, California, USA
- <sup>9</sup>Pediatrics / Pediatric Neurology, Mattel Children's Hospital UCLA, Los Angeles, California, USA
- <sup>10</sup>Psychology, University of Calgary, Calgary, Alberta, Canada
- <sup>11</sup>Physiotherapy Department, University Hospitals Dorset NHS Foundation Trust, Poole, UK
- <sup>12</sup>The FA Centre for Para Football Research, The Football Association, Burton-Upon-Trent, UK
- <sup>13</sup>Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation/Harvard Medical School, Boston, Massachusetts, USA
- <sup>14</sup>The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA
- <sup>15</sup>Orthopedics and Sports Medicine, Pediatrics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA
- <sup>16</sup>Department of Radiology, University of Calgary, Calgary, Alberta, Canada
- <sup>17</sup>Children's Hospital of Eastern Ontario Research Institute, Ottawa, Ontario, Canada
- <sup>18</sup>Psychology, University of Missouri Kansas City, Kansas City, Missouri, USA
- <sup>19</sup>Psychological and Neurobehavioral Associates, Inc, Miami, Florida, USA
- <sup>20</sup>Rehabilitation, Laval University, Quebec, Quebec, Canada
- <sup>21</sup>School of Health and Related Research, University of Sheffield, Sheffield, UK
- <sup>22</sup>Family Medicine, University of Washington, Seattle, Washington, USA
- <sup>23</sup>Departments of Rehabilitation Medicine, Orthopaedics and Sports Medicine and Neurological Surgery, University of Washington, Seattle, Washington, USA
- <sup>24</sup>University of Calgary, Calgary, Alberta, Canada
- <sup>25</sup>Institute for Sport Exercise and Health, University College Hospital London, London, UK
- <sup>26</sup>Florey Institute of Neuroscience and Mental Health - Austin Campus, Heidelberg, Victoria, Australia
- <sup>27</sup>La Trobe Sport and Exercise Medicine Research Centre, Melbourne, Victoria, Australia
- <sup>28</sup>Neurosurgery, Medical College of Wisconsin, Milwaukee, Wisconsin, USA
- <sup>29</sup>Sports Medicine, Children's Hospital Boston, Boston, Massachusetts, USA
- <sup>30</sup>Emergency Medicine, Children's Hospital Boston, Boston, Massachusetts, USA
- <sup>31</sup>Musculoskeletal Medicine, Royal Orthopaedic Hospital, Birmingham, UK
- <sup>32</sup>Libraries, University of Victoria, Victoria, British Columbia, Canada
- <sup>33</sup>Major League Soccer, New York, New York, USA
- <sup>34</sup>Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada
- <sup>35</sup>International Concussion and Head Injury Research Foundation, London, UK
- <sup>36</sup>University of Calgary Faculty of Kinesiology, Calgary, Alberta, Canada
- <sup>37</sup>Loughborough University, Loughborough, UK

<sup>38</sup>Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Calgary, Alberta, Canada

**Twitter** Jon S Patricios @jonpatricios, Vicki Anderson @VickiAn28323584, Christopher C Giza @griz1, Osman Hassan Ahmed @osmanhahmed, Christina L Master @drtinamaster, Pierre Fremont @pfremo, Kimberly G Harmon @DrKimHarmon, Zahra Premji @ZapTheLibrarian, Kenzie Vaandering @kenzievaan, Nick Webborn @SportswiseUK and Kathryn J Schneider @Kat\_Schneider7

**Contributors** GAD served as the primary author and responsible for all aspects of the project, including initial preparation, coordination, review, editing and final preparation of the Child SCOAT6 tool. JSP served as the primary author of the systematic review and development of the SCOAT manuscript and tool. All co-authors contributed to the development and critical review of the Child SCOAT6 tool, and approved the final version of the tool.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** GAD is a member of the Scientific Committee of the 6th International Consensus Conference on Concussion in Sport; an honorary member of the AFL Concussion Scientific Committee; Section Editor, Sport and Rehabilitation, NEUROSURGERY; and has attended meetings organised by sporting organisations including the NFL, NRL, IIHF, IOC and FIFA; however has not received any payment, research funding, or other monies from these groups other than for travel costs. Dr JSP, Editor BJSM (honorary), Member of World Rugby Concussion Advisory Group (unpaid), Independent Concussion Consultant for World Rugby (fee per consultation), Medical consultant to South African Rugby (unpaid), Co-chair of the Scientific Committee, 6th International Conference on Concussion in Sport (unpaid), Board member of the Concussion in Sport Group (unpaid), Scientific Board member, EyeGuideTM (unpaid). Dr LP CASEM Board Member, President-Elect 2022-2023NIH R34 Grant for EPICC Study (Eye Problems in Concussed Children), Site PI Speaker at various conferences. Dr VA Financial: Australian National Health and Medical Research Council and Medical Research Future fund: research grants. Royalties: Pearson Publishing (Test of Everyday Attention) Collaboration: Australian Football League (Partnership agreement to fund research – funds go to my institute). Boards: Editorship: Journal of Neuropsychology, Neuropsychology, Journal of Clinical NIH NINDS (R01 NS110757 2019-2024); NINDS(U54 NS121688 2021-2026); UCLA Brain Injury Research Center, UCLA Steve Tisch Brain SPORT program, Easton Clinic for Brain Health Clinical Consultant (provide clinical care to athletes): NBA, NFL-Neurological Care Program, NHL/NHLPA, Los Angeles Lakers Advisory Board (Non compensated): Major League Soccer, National Basketball Association, US Soccer Federation. Advisory Board (Compensated): Highmark Interactive MedicoLegal: One or two cases annually Speaker's Bureau: None. Stock Shareholder: Highmark Interactive stock options (2018). Other Financial or Material Support: Book royalties – Blackwell/Wiley Publishing: Prioritized Neurological Differential Diagnosis Other: None. Dr KOY: is Editor-in-Chief of the journal Neuropsychology and receive an editorial stipend from the American Psychological Association. I am an unpaid consulting editor for the journals Archives of Clinical Neuropsychology and Journal of Head Trauma Rehabilitation. I am an unpaid member of the Scientific Advisory Committee for Brain Injury Canada. I am the chair of the Canadian Concussion Network, which is funded by a grant from Canadian Institutes of Health Research (CIHR) to my institution; I am principal

applicant on the grant but receive no income from it. I am a principal investigator on another grant from CIHR from which I derive no income. I am a co-investigator on research grants from CIHR, the US National Institutes of Health (NIH), Brain Canada Foundation and National Football League Scientific Advisory Board; I derive income only from the grant from NIH. I serve as a member of a CIHR grant review panel for which I receive a small honorarium. I receive book royalties from Guilford Press and Cambridge University Press. I have received travel support and honorarium for presentations to multiple organisations. I served or serve on the following committees/boards for which I receive(d) honorarium: 1. Independent Data Monitoring Committee (IDMC), Care for Post-Concussive Symptoms Effectiveness (CARE4PCS-2) Trial, National Institute for Child Health and Human Development 2. Observational Study Monitoring Board (OSMB), Approaches and Decisions in Acute PediatricTBI (ADAPT) Trial, National Institute of Neurological Disorders and Stroke National Research Advisory Council, National Pediatric Rehabilitation Resource Center, Center for Pediatric Rehabilitation: Growing Research, Education and Sharing Science (C-PROGRESS), Virginia Tech University. Dr OHA is a Senior Physiotherapist at University Hospitals Dorset NHS Foundation Trust (England) and is Para Football Physiotherapy Lead/Para Football Classification Lead at the Football Association (England). He also works on a consultancy basis with the Football Association as the squad physiotherapist to the England Cerebral Palsy Football squad and teaches on the Football Association's Advanced Trauma and Medical Management in Football course on a consultancy basis. He has a Visiting Senior Lecturer position at the University of Portsmouth, England (unpaid). He sits on several disability sport committees including Para Football Foundation as Medical Unit Co-Lead (unpaid), the International Federation of Cerebral Palsy Football as Medical and Sports Science Director (unpaid) and the International Blind Sports Association as Medical Committee member (unpaid). He has Associate Editor positions at the British Journal of Sports Medicine (unpaid) and BMJ Open Sports my time is supported by National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award number K23NS128275-01Dr. Christina Master reports no financial COI Volunteer positions: Concussion team physician, Shipley School Board of Trustees, American College of Sports Medicine Board of Directors, American Medical Society for Sports Medicine Board of Directors, Pediatric Research in Sports Medicine Executive Committee, Council on Sports Medicine and Fitness, American Academy of Pediatrics Advisory Board, Untold Foundation, Pink Concussions, Headway Foundation Editorial Board, Journal of Adolescent Health, Frontiers in Neuroergonomics, Exercise, Sport, and Movement. Dr Geoff Schneider is an owner of a multidisciplinary practice (managing patients with MSK pain disorders). He is a board member of Hockey Calgary (Calgary, AB, Canada) and Chair of the Alberta Association of Physiotherapy. He received funding for the administrative aspects of the writing of two of the systematic reviews that informed the consensus process. Dr JVI is the founder of R2P Concussion Management. Dr RJE is a paid consultant for the NHL and cochair of the NHL/NHLPA Concussion Subcommittee. He is also a paid consultant and chair of the Major League Soccer concussion committee and a consultant to the US Soccer Federation. He previously served as a neuropsychology consultant to Princeton University Athletic Medicine and Eye Guide. He is currently a co-PI for a grant funded by the NFL (NFL-Long) through Boston Children's Hospital. He occasionally provides expert testimony in matters related to MTBI and sports concussion and occasionally receives honoraria and travel support/reimbursement for professional meetings. Dr PF is a co-investigator on

a research grant from the NFL's "Play Smart. PlaySafe." Initiative and an Executive committee member of the Canadian Concussion Network (financed by the Canadian Institute of Health Research). He received honorarium for an Expertgroup discussion on blood biomarkers for concussion in December 2020. Dr GWF has received travel expenses to attend academic meetings from World Rugby. He has also collaborated on research projects with World Rugby as chief or co-investigator. He is previous associate editor of the British Journal of Sports Medicine. He has not received any other payments or support from any sporting or commercial bodies. He has no other conflicts of interest. Dr Stanley A Herring Co-founder and senior advisor, The Sports Institute at UW Medicine (unpaid) Centers for Disease Control and Prevention and National Center for Injury Prevention and Control Board Pediatric Mild Traumatic Brain Injury Guideline Workgroup (unpaid) CISC (travel support) NCAA Concussion Safety Advisory Group (unpaid) Team Physician, Seattle Mariners Former Team Physician, Seattle Seahawks occasional payment for expert testimony travel support for professional meetings. Dr KGH Research Development Director, Pac-12 Conference Member, Pac-12 Brain Trauma Task Force Member, NFL Head Neck and Spine Committee Deputy Editor, British Journal of Sports Medicine Head Football Physician, University of Washington. Ms Holte received an honorarium for administrative aspects of the reviews. Dr ML is the CMO GB Boxing, CMO GB Snowsports. NE Director GB Taekwondo. NE Director SWA (share options). Director Active Movement. Director GB Obstacle course racing. Founder and medical board member of Safe MMA. Director of Marylebone Health Group. Private medical practice at ISEH 170 Tottenham Court Road. Private medical practice Marylebone Health Group. Dr MM Sport and exercise medicine physician working in private consulting practice. Shareholder of Olympic Park Sports Medicine Centre in Melbourne. Ex-senior physician at the Hawthorn Football Club (AFL) Ex-Chief Executive Officer of the AFL Doctors Association. Research grants received from the Australian Football League, outside the submitted work. Travel support received from the Australian Football League, FIFA and the International Olympic Committee to attend and present at international conferences. Member of the Scientific Committee for the 6th International Consensus Conference on Concussion in Sport. Honorary member of the International Concussion in Sport Group. Honorary member of the Australian Rugby Union Concussion Advisory Group. Independent Concussion Consultant for World Rugby. Dr MM has received research funding to the Medical College of Wisconsin from the National Institutes of Health, Department of Veterans Affairs, Centers for Disease Control and Prevention, Department of Defense, National Collegiate Athletic Association, National Football League and Abbott Laboratories. He receives book royalties from Oxford University Press. He serves as clinical consultant to Milwaukee Bucks, Milwaukee Brewers, and Green Bay Packers and is Co-Director of the NFL Neuropsychology Consultants without compensation. He serves as consultant for Neurotrauma Sciences, Inc. He receives travel support and speaker honorariums for professional activities. WPM - I receive royalties from ABC-Clio publishing for the sale of the books, Kids, Sports and Concussion: A guide for coaches and parents and Concussions; from Springer International

for the book Head and Neck Injuries in Young Athlete; and from Wolters Kluwer for working as an author for UpTo Date. My research is funded, in part, by philanthropic support from the National Hockey League Alumni Association through the CCG Pro-Am Tournament and a grant from a grant from the National Football League. POH - is an employee of Marker Diagnostics UK Ltd. He has worked on a sessional basis for the English RFU and FA. He has previously received research funding from the English RFU, ECB and The Drake Foundation. Dr ZP No COI to declare. Dr MP declares the following: Consultant, CMO, Major League Soccer, Senior Advisor, NFL Head, Neck NCAA-CARE-DoD 2.0, ended 2020. Have received honoraria and reimbursement for travel for speaking and conferences attended. Have written chapters for UpToDate and received royalties for the Netter's Sports Medicine textbook. Have provided work as an expert for cases involving concussion, team physician and other sports medicine topics. IJS - does not have any conflicts of interest to disclose. Dr MT is employed full-time as the CEO and Medical Director of ICHIRF - a paid post he has held since April 2015. Hon Medical Adviser to the Professional Riders Insurance Scheme (PRIS) - discretionary honorarium Member of the Premier League Head Injury Advisory Group (HIAG) - no remuneration Director of ICHIRF Ireland Ltd - no remuneration Honorary Medical Adviser to the Concussion Foundation Ltd - no remuneration Member of the expert panel for the Dept of Digital, Culture, Media and Sport review into concussion in amateur sport - no remuneration. Attendance at conferences or meetings as a guest speaker - reimbursement of travel expenses, complimentary registration and payment of hotel accommodation and meals by the organising committee. No stocks or options in any concussion-related company. No consultancies, board or editorial positions related to concussion. KV - has no conflicts of interest to disclose. Dr NW Chair, British Paralympic Association (voluntary) IPC Medical Committee Member (voluntary) Concussion in Para Sports (CIPS), founding member (voluntary) BJSM Editorial Board member (voluntary) Sports Horizon, Board of Directors - equity share - see <https://www.sportshorizon.co.uk>. KJS has received grant funding from the Canadian Institutes of Health Research (CIHR), NFL Scientific Advisory Board, International Olympic Committee Medical and Scientific Research Fund, World Rugby, Mitacs Accelerate, University of Calgary, with funds paid to her institution and not to her personally. She is an Associate Editor of BJSM (unpaid), Independent consultant to World Rugby and has received travel and accommodation support for meetings where she has presented. She coordinated the writing of the systematic reviews that informed Amsterdam International Consensus on Concussion in Sport, for which she has received an educational grant to assist with the administrative costs associated with the writing of the reviews (with funds paid to her institution). She is a member of the AFL Concussion Scientific Committee (unpaid position), Brain Canada (unpaid positions) and Board member of the Concussion in Sport Group (CISC)(unpaid). She works as a physiotherapy consultant and treats athletes of all levels of sport from grass roots to professional.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2023-106984>).



**To cite** Davis GA, Patricios JS, Purcell LK, *et al*. *Br J Sports Med* 2023;**57**:672–688.

Accepted 5 June 2023

*Br J Sports Med* 2023;**57**:672–688.  
doi:10.1136/bjsports-2023-106984

#### ORCID iDs

Gavin A Davis <http://orcid.org/0000-0001-8293-4496>  
Jon S Patricios <http://orcid.org/0000-0002-6829-4098>  
Keith Owen Yeates <http://orcid.org/0000-0001-7680-2892>  
Osman Hassan Ahmed <http://orcid.org/0000-0002-1439-0076>  
Cheri Blauwet <http://orcid.org/0000-0001-8568-1009>  
Christina L Master <http://orcid.org/0000-0002-6717-4270>  
Geoff Schneider <http://orcid.org/0000-0002-4485-1769>  
Jacqueline van Ierssel <http://orcid.org/0000-0001-5519-8526>  
Ruben J Echemendia <http://orcid.org/0000-0001-6116-8462>  
Pierre Fremont <http://orcid.org/0000-0003-2810-8382>  
Gordon Ward Fuller <http://orcid.org/0000-0001-8532-3500>  
Kimberly G Harmon <http://orcid.org/0000-0002-3670-6609>  
Mike Loosemore <http://orcid.org/0000-0002-4855-0744>  
Michael McCrea <http://orcid.org/0000-0001-9791-9475>  
Patrick O'Halloran <http://orcid.org/0000-0002-1185-3485>  
Zahra Premji <http://orcid.org/0000-0002-6899-0528>  
Michael Turner <http://orcid.org/0000-0003-2323-2456>  
Kenzie Vaandering <http://orcid.org/0000-0002-2342-5373>  
Nick Webborn <http://orcid.org/0000-0003-3636-5557>  
Kathryn J Schneider <http://orcid.org/0000-0002-5951-5899>